

PREOP CONTINUATION OF MEDICATIONS – continue **all** meds except:

ANTICOAGULANTS (including aspirin): Instructions on holding anticoagulants need to come from the prescriber, the surgeon, or a consultant

NSAIDs: Hold for 4 days (unless otherwise instructed by the surgeon)

INSULINS: Continue as normal until the morning of procedure:

- If BG > 140, take one-half of AM dose of long-acting insulins (LANTUS, LEVEMIR, TOUJEO, TRESIBA)
- Pumps – suggest decreasing the basal rate by one-half (patient may override this instruction)
- Hold all other insulins

ORAL DIABETIC MEDS: Hold the day of surgery

- If on SGLT2 inhibitors (FARXIGA, INVOKANA, JARDIANCE, BRENZAVVY) and having major surgery, notify anesthesia

INJECTABLE NON-INSULIN DIABETIC MEDS: For those on once daily dosing, just hold the day of surgery. For those on weekly dosing, hold for 7 days whenever possible

DIURETICS: Hold the day of surgery if no history of HF

HERBALS AND VITAMIN E:

- Ideally hold for 7 days (will not cancel case if this is not possible)
- Cannabis in all forms – Do not consume two hours before arrival

Type-A MAO Inhibitors for Depression: Hold for 7 days, but only under psychiatric supervision
MARPLAN, NARDIL, PARNATE

For Bariatric Patients Only: Hold ACE inhibitors and ARBs for 24 hours

NARCOTIC LIMITERS: Can continue as normal for minor surgery, otherwise consult anesthesia for instructions

Generic Name	Trade Name	Dosing	Frequency	Use
Buprenorphine	BELBUCA	ODT/SL	Daily	Narc control
	BRIDAXI 8-32	ODT/SL	Weekly	Narc control
	BRIDAXI 64-128	SQ	Monthly	Narc control
	BUTRANS	Patch	Weekly	Narc control
	SUBLOCADE	SQ	Monthly	Narc control
Buprenorphine + naloxone	SUBOXONE	ODT/SL	Daily	Narc control
	ZUBSOLV	ODT/SL	Daily	Narc control
Naltrexone	REVIA	PO	Daily	Narc control
	VIVITROL	IM	Monthly	Alcoholism
Naltrexone with antidepressant	CONTRAVE	PO	Daily	Weight loss